
MAYNE Is. MAP I of 3: Distribution of Shoreline Types The Mayne Island shoreline is largely bedrock and resistant to erosion. There are several sand and shell pocket beaches that are very important recreationally. The north side of Mayne Island is exposed to northerly storm winds and waves. The west and south sides of Mayne Island are largely protected from wind and waves by Galiano Island and North Pender Island respectively. There are a few areas of tidal sand or mudflats, and several areas of saltmarsh, all of which are sensitive to human disturbance. There are high current areas in the passes around the island, particularly Active Pass to the west. Drainage Conveyance Zones (wetlands, creeks & moist soil areas) Watershed Boundaries low rock/boulder marsh/fine sediment boulder / cobble structurally altered* **MAYNE ISLAND:** I.7 Km (4%) of Structurally Altered pebble / sand Shoreline



SEA CLIFF rocky shore with

steep slopes

(7.3 Km)

• 17% of shoreline

LOW ROCK/BOULDER

• moderate to high rocky shore with slopes of sediment low slopes (often eroding) 66% of shoreline

 Mayne Island has no (27.7 Km)

BOULDER/COBBLE

I Km

BLUFF

bluff shores

• boulder - cobble cover on beach (often indicates eroding shoreline)

• 4% of shoreline (1.7 km)

PEBBLE/SAND

stable or accreting pebble-sand (or shell) beaches (may be eroding where sediment supply is interrupted).

• 10% of shoreline (4.2 Km)

MARSH/FINE SEDIMENT

DATE: October 25, 2011

• low energy shorelines with sediment inputs from watersheds nearby

• 2% of shoreline (9.2 Km)

Rock (Hard) Shorelines





Sediment (Soft) Shorelines






Refer to Islands Trust Report "Gulf Islands Shoreline Mapping - Project Methodology" July, 2011 for detailed map informatio



ISLAND ENVIRONMENTS are shaped by two primary or formative systems: 1) watershed systems; and 2) longshore systems.

Watersheds are driven by runoff, and longshore systems are driven by waves and ocean currents. Any attempt to understand the islands, including discussions about land use planning, must be framed by these systems. Within this framework all other systems (natural systems like forests, wetlands, eelgrass beds etc, and human systems like roads, buildings, etc) are organized and structured.

MAYNE ISLAND belongs to a class of sea coast know as sheltered shoreline because it is not exposed to the open sea. Nevertheless, wind wave and current activity remains the controlling force along the island's perimeter. The strength of this system, however, varies appreciably depending on the orientation and form of the shoreline, near shore water depth, and other factors. For example, headlands facing storm waves are subject to the greatest wind and wave force, whereas bays and estuaries are subject to the least. Not surprisingly,



Sediment Movement:

Accretion Shorelines (adding sediment) Erosion Shorelines (losing sediment)

Watershed Sediment Inputs to Shoreline System



Localized Sediment Movement Direction (Small Scale)





Predominant Direction of Wave Energy

Making Sense of the Energy Systems:

The coloured shorelines on this map indicate that for Mayne Island, the highest wave exposures occur on the north side of the island. The arrows indicate that the predominant wave energy system flow direction is from south to north.

This seemingly conflicting information can be explained by examining the difference between wave exposure and prevailing storms. Wave exposure is a function of wave fetch (as described in the inset to the right) and wind strength from a given direction. The north shore of Mayne Island has a long fetch relative to other Mayne Island shorelines, and our region receives periodic outflow winds from the north – this combination of factors results in the north facing beaches having a high wave exposure rating. Predominant energy flow direction, on the other hand, is the cumulative effect of storms over time. Our region receives some winter outflow winds from the north, but the majority of strong winds and storms come from the southeast. For Mayne Island, this means that the dominant sediment movement (relatively little sediment exists on the Mayne Shoreline because it is predominantly rocky) direction is northward, driven by the south-easterly storms.






DATE: October 25, 2011

Accretion Shorelines:

Sediment accumulation (accretion) is typically associated with lower energy environments along the shorelines.

Accretion features include sandy beaches, beach berms, pocket beaches or storm berms, and are often high value recreation features or wildlife habitats.



Erosion Shorelines:

Eroding shorelines are typically associated with higher energy environments along the shorelines, like headlands, high exposure sediment shorelines or points of land.

Eroding shorelines feed the sediment transport system and halting erosion can have severe impacts on the shoreline sediment movement system and 'downstream' beaches. Adequate setbacks for buildings and facilities are critical.



Trees and vegetation damaged or shaped by the wind along shorelines are good indicators of high wind exposure.

siting buildings and facilities in these locations to ensure they are adequately set back from the shoreline.

Caution should be exercised when

Wave Exposure & the Sediment System



PROJECT FUNDING GENEROUSLY PROVIDED BY:







Refer to Islands Trust Report "Gulf Islands Shoreline Mapping - Project Methodology" July, 2011 for detailed map informatic



MAYNE Is. MAP 3 of 3: Shoreline Values & Vulnerability This map is intended to give a general impression of areas along the Mayne Island shoreline that are considered valued and/or vulnerable to change. Value refers to areas or features of high ecological or recreational significance. Vulnerability refers to: 1) Natural areas or features vulnerable to human disturbance; or 2) Buildings or facilities, vulnerable to disturbance from natural or human-altered system processes. It is important to note that the various features highlighted on this map are in many cases both vulnerable in some way, and valued. Saltmarshes for example are of high ecological value, and are also vulnerable to accumulation of pollutants potentially contained in island runoff. Watershed Boundaries. Drainage Conveyance Zones-(wetlands, creeks & moist soil areas) Watershed Areas Draining to Sensitive Shorelines (i.e. Very **COMMUNITY INPUT** Protected Waters and Salt Marsh) THIS IS YOUR SPACE - tell us what we have missed or where you think the mappers have erred to help us build FACILITY SITING & SETBACKS a more comprehensive values and vulnerability resource! Feel free to make notes right on the map also. Riparian vegetation helpsf building setback from shore protects stabilize shoreline and reduces vulnerability of recreational values structures to shoreline Sediment movement septic systems less constrained and fields along beaches more effective uninterrupted Lower Facility Risk - Improved Shore Protection

small setback building and seawall reduce recreational necessitates seawall access to beach protection seawall interrupts sediment movement along shoreline septic systems too close to shore keep soils moist and prone to erosion Higher Facility Risk - Loss of Shore Values

DATE: October 25, 2011 Refer to Islands Trust Report "Gulf Islands Shoreline Mapping - Project Methodology" July, 2011 for detailed map information

LOW LYING AREAS

• Areas 0-4 m in elevation above existing Mean Sea Level and greater than 50 m of width from the shore are mapped as 'vulnerable' to sea level rise.

Current BC Provincial Government guidelines suggest up to I metre of sea level rise over the next 100 years (www.env.gov.bc.ca). Sea level rise may cause increased shoreline vulnerability to land-based activities by causing such effects as increased flooding in low lying areas or softening of sediment shorelines and increased shoreline erosion. These effects could be further exacerbated by storm surges and changing climatic conditions.

SOFT SHORELINES

• Sediment shorelines are typically associated with high recreational values and high ecological values (pocket beaches, estuaries, etc).

VERY PROTECTED WATERS

- Shorelines highly protected from wave exposure
- Poor water circulation increases vulnerability to water pollution





Areas of High Ecological or Recreational Significance



PROJECT FUNDING GENEROUSLY PROVIDED BY:







