


# Sensitive Ecosystems on North Pender Island Airphoto - 2004

Scale: 1:10,000  
UTM Projection Zone 10 NAD83, Contour Interval 10 metres  
February 2007

Ecosystem composition is complex and often contains a dominant ecosystem with secondary and tertiary ecosystems. In this map the dominant ecosystem has a solid shading and the secondary and tertiary ecosystems are identified by hatched lines.

Example of dominant Woodland ecosystem with a secondary Herbaceous ecosystem  
Plotted Feb 15 2007



Sensitive Ecosystems	
Sensitive ecosystems are fragile and/or rare, or are ecologically important because of the diversity of species they support.	
<b>Old Forest (OF):</b>	Old forest ecosystems are structurally complex stands composed mainly of shade-tolerant and regenerating tree species in late successional stages. They are generally 70 years or more. The inventory can include large and open-woody areas in stages of decomposition, and a fully developed moss layer.
<b>Importance:</b>	Due to the lack of disturbance, old forest ecosystems are often associated with rich communities of plants and animals that may be dependent upon the unique environmental conditions created by these forests.
<b>Subclasses:</b>	<ul style="list-style-type: none"> <li>of (old forest) - forests dominated by large, old broadleaf trees</li> <li>mf (mixed conifer) - forests dominated with a mixture of coniferous and broadleaf trees (&lt;75% coniferous and &gt;25% broadleaf)</li> <li>cc (coniferous) - conifer-dominated (&gt;75%) forests generally &gt;250 years</li> </ul>
<b>Woodland (WD):</b>	Definition: of pure or mixed stands of Garry oak or mixed stand of arbutus and Douglas fir. Mature big leaf maple can also be found in sites designated as woodland.
<b>Importance:</b>	Woodlands are nationally, provincially and regionally rare and highly fragmented. A rich assemblage of plants, insects, reptiles and birds are often found in these ecosystems due to the food resources, habitat and proximity to the ocean. Garry oak woodlands, for example support the highest plant species diversity of any terrestrial ecosystem in British Columbia and are especially vulnerable to rural development.
<b>Subclasses:</b>	<ul style="list-style-type: none"> <li>bd (broadleaf) - dominant broadleaf with &lt;15% coniferous species</li> <li>mc (mixed conifer and deciduous) - mixed conifer and broadleaf with a minimum of 25% cover of either group is included in the total tree cover</li> </ul>
<b>Herbaceous (HB):</b>	Definition: Non-forested ecosystems (less than 10% tree cover), generally with shallow soils. They include bedrock outcroppings, large openings within forested areas, sph. dunes and shorelines vegetated with grasses and herbs.
<b>Importance:</b>	Terrestrial Herbaceous ecosystems are characterized by thin soils which are easily disturbed. Herbaceous plants can be easily trampled or displaced into bare rock where they cannot reestablish. They are highly vulnerable to a range of human disturbance factors including residential development and various recreational uses.
<b>Subclasses:</b>	<ul style="list-style-type: none"> <li>hb (herbaceous) - non-forested, less than 10% tree cover, generally shallow soils, often with exposed bedrock, predominantly a mix of grasses and forbs, also lichens and mosses</li> <li>ca (coastal herbaceous) - mostly dunes or wet, influenced by the marine environment and characterized by less than 20% vegetation cover of grasses, forbs, mosses and lichens</li> <li>sp (spit) - higher elevation of beach, comprised of sand or gravel deposited by longshore drifting, low to moderate cover of salt-tolerant grasses and herbs</li> <li>du (dunes) - ridge or hill, or beach area created by windblown sand; may be more or less vegetated depending on depositional activity, beach dunes will have low cover of salt-tolerant grasses and herbs</li> <li>sh (shrub) - &lt;20% of total vegetation cover is shrub cover, with grasses and herbs</li> <li>ro (rock) - rock outcrop not dominated by shrubs</li> </ul>
<b>Riparian (RI):</b>	Definition: Areas adjacent to water bodies (rivers, lakes, ocean, wetlands) which are influenced by factors such as erosion, sedimentation, fluctuating water tables, tidal influences or poor drainage conditions.
<b>Importance:</b>	Riparian ecosystems support a disproportionately high number of vascular plant, moss, amphibian and small mammal species for the area they occupy.
<b>Subclasses:</b>	<ul style="list-style-type: none"> <li>fl (low bench floodplain) - flooded at least every other year for moderate periods of growing season; plant species adapted to extended flooding and erosion; low or tall shrubs most common</li> <li>fm (medium bench floodplain) - flooded every 1-6 years for short periods (10-25 days); deciduous or mixed forest dominated by species tolerant of flooding and periodic sedimentation; trees occur on elevated microsites</li> <li>fr (high bench floodplain) - only periodically and briefly inundated by high waters, but lengthy subsurface flow in the rooting zone; typically conifer-dominated floodplains of larger coastal rivers</li> <li>tr (trough) - narrow linear communities along with open water bodies (rivers, lakes and ponds) where there is no floodplain; irregular flooding</li> <li>gr (gully riparian) - watercourse is within a steep sided V-shaped gully</li> <li>ri (river) - watercourse is large enough to represent &gt;10% of the polygon</li> <li>st (strath) - strath-dominated floodplain or savanna</li> </ul>
<b>Wetland (WN):</b>	Definition: Areas that are saturated or inundated with water for long enough periods of time to develop vegetation. This may result from flooding, fluctuating water tables, tidal influences or poor drainage conditions.
<b>Importance:</b>	Wetland ecosystems are sensitive and important because they exhibit rarity, high biodiversity, fragility, specialized habitat, specialized functions and connectivity.
<b>Subclasses:</b>	<ul style="list-style-type: none"> <li>bg (bog) - nutrient poor wetland, on organic soils (sphagnum peat), water source predominantly from precipitation; may be tree or shrub dominated</li> <li>fw (fen) - nutrient medium wetland (sedge peat) where ground water inflow is the dominant water source; open water channels common; dominated by sedges, grasses and mosses</li> <li>me (marsh) - wetland with fluctuating water tables, often with shallow surface water; usually organically enriched mineral soils; dominated by rushes, reeds, grasses and sedges</li> <li>sw (swamp) - poor to very rich wetland on mineral soils or with an organic layer over mineral soil, with gently flowing or seasonally flooding water table; woody vegetation</li> <li>sp (shallow water) - standing or flowing water less than 2m deep; transition between deep water bodies and other wetland ecosystems (i.e. bogs, swamps, fens, etc.); often with vegetation rooted below the water surface</li> <li>wt (wet meadow) - periodically saturated but not inundated with water; organically enriched mineral soils; grasses, sedges, rushes and forbs dominate</li> </ul>
<b>Cliff (CL):</b>	Definition: Very steep slope, often exposed bedrock, may include steep-sided sand dunes.
<b>Importance:</b>	Open ledges and horizontal features on cliffs are known to provide nesting sites. Cliff crevices are used for roosting bats while deep crevices are used for shelter and overwintering of snakes and lizards.
<b>Subclasses:</b>	<ul style="list-style-type: none"> <li>cc (coastal cliffs) - cliffs with a marine influence, generally near vertical bedrock with accumulation of soil limited to fissures and ledges</li> <li>ic (inland cliffs) - inland cliffs typically formed as a result of erosion, catastrophic failures or mass wasting. Generally characterized by rapid drainage and the accumulation of soil that is limited to bedrock fissures and ledges</li> </ul>
<b>Freshwater (FW):</b>	Definition: Freshwater ecosystem includes bodies of water such as lakes and ponds that usually lack floating vegetation.
<b>Importance:</b>	Freshwater ecosystems are home to numerous organisms such as fish, amphibians, aquatic plants, and invertebrates.
<b>Subclasses:</b>	Lakes and ponds play a vital role in the life cycle of many species.
<b>la (lake)</b>	a naturally occurring static body of water, greater than 2m deep in some portion
<b>pd (pond)</b>	a small body of water greater than 2m deep, but not large enough to be classified as a lake
<b>Intertidal (IT):</b>	Definition: Areas that link the marine and terrestrial environments and generally consist of mudflats and beaches.
<b>Importance:</b>	Intertidal ecosystems are one of the most biologically diverse natural communities because it is where the land and water meet. The many shoreline habitats provide wildlife with food, den and burrow sites, nursery areas, flyways and travel corridors, perching and sunning sites and protection from weather and predators.
<b>Subclasses:</b>	<ul style="list-style-type: none"> <li>me (mudflat) - flat, plain-like areas dominated by fine-textured sediments and exposed at low tide; includes estuaries</li> <li>be (beach) - areas that express sorted sediments, reworked by wave action in recent times</li> </ul>
<b>Other Important Ecosystems</b>	Other important ecosystems have high biodiversity values.
<b>Mature Forest (MF):</b>	Definition: Usually conifer-dominated, occasionally deciduous, dry to moist forest types; structural stage 6, generally >80yr.
<b>Importance:</b>	Within 20 years, many Mature Forests that were logged early in this century will become Old Forests. The biodiversity values of Mature Forests generally become higher with age. This means it will be difficult to sustain more and larger species of plants and animals.
<b>Landscaping connectivity:</b>	Mature Forest stands provide connections between other natural areas that promote the movement and dispersal of many forest-dwelling species across the landscape.
<b>Buffer:</b>	Mature Forest can minimize disturbance to sensitive ecosystems that occur within or adjacent to the forest patch. When they border or surround wetlands, patches of old forest or other sensitive ecosystems, the Mature Forest area serves an important role in buffering the adjacent sensitive areas.
<b>Subclasses:</b>	<ul style="list-style-type: none"> <li>cc (conifer dominated) - greater than 75% coniferous species</li> <li>mc (mixed conifer and deciduous) - a minimum of 25% cover of other group is included in the total tree cover</li> </ul>
<b>Other Mapped Ecosystems</b>	
<b>Young Forest (YF):</b>	On this map, Young Forest ecosystems are depicted ONLY when they occur in a mosaic with other sensitive ecosystems. For example, a Young Forest dominates the polygon but there is also a distinct wetland within the polygon boundary.

### What is a Sensitive Ecosystem?

For the purpose of this study, an ecosystem is considered to be a portion of the landscape with relatively uniform dominant vegetation. Sensitive ecosystems are those which are fragile and/or rare, or those ecosystems which are ecologically important because of the diversity of species they support.

### Rationale

Intense development pressure fueled by population and economic growth has fragmented and degraded many terrestrial ecosystems. A high proportion of these ecosystems are now designated as 'at risk' in BC. Sensitive ecosystems typically have high biological diversity and are a vital part of the landscape. They provide ecosystem services for a healthy economy and for social well being. They regulate climate, clean water, generate and clean soils, recycle nutrients and pollinate our crops. To protect these areas, sensitive ecosystems must be located, identified and mapped. From 1993 to 1999 the Provincial and Federal Governments completed a Sensitive Ecosystems Inventory of East Vancouver Island and the Gulf Islands. This mapping product is an updated version of that product.

### Purpose

The purpose of this Sensitive Ecosystems map is to identify the location of sensitive ecosystems. The goal of this mapping exercise is to encourage informed land use decisions that will conserve sensitive ecosystems. This map and the accompanying data provide site-specific ecological information that can be used to flag sites of conservation concern, to promote land stewardship and to prompt detailed field surveys and consideration of ecological values before changes to the land are initiated.

### Methodology

Mapping methods are based on the Resource Information Standards Committee (RISC) Standard for Terrestrial Ecosystem Mapping (TEM) in BC. This Sensitive Ecosystems map was themed from TEM data using the RISC Standard for Mapping Ecosystems at Risk in BC. Field survey protocols followed Describing Terrestrial Ecosystems in the Field (RISC 1998).

### Data Limitations

The Sensitive Ecosystems map is a tool to alert decision makers to the existence of sensitive ecosystems. However, when land-use changes are proposed, detailed on-the-ground site assessments are necessary. For sites that were not field checked, the accuracy of the data depends heavily on the expertise, local knowledge, and professional judgment of the mapper and the quality and quantity of available source data. Because the area is changing rapidly, reference to the data set(s) used as the information source is advised.

### What can be done to protect the sensitive ecosystems?

- Retaining or creating vegetated buffers around sensitive ecosystems to isolate them from outside disturbances;
- Controlling land and water access to fragile ecosystems;
- Controlling invasive species;
- Allowing natural disturbances to occur;
- Maintaining water quality

### If development must occur, develop carefully!

Conduct an ecological inventory to identify the existing flora and fauna and to locate any threatened or endangered plant and animal species, plant communities, and habitat features needing protection.

Plan and implement all development activities in a manner that will not adversely affect or disturb the sensitive ecosystem. Consult a qualified professional to interpret the ecological inventory data and work to incorporate designs that maintain the functions and values of the natural ecosystem.

### Acknowledgements:

- Project Co-ordinator: Ardice Todoshchuk, Islands Trust Fund/Islands Trust
- Terrestrial Ecosystem Mapping: Andy MacKinnon - BC Ministry of Forests and Range
- Todd Golumbia - Gulf Islands National Park Reserve
- Tara Sharma - Gulf Islands National Park Reserve
- Sensitive Ecosystem Mapping Conversion Tables: Carmen Cadrin - BC Ministry of Environment
- Jo-Anne Stacey - BC Ministry of Environment
- Andy MacKinnon - BC Ministry of Forests and Range
- GIS Mapping Support: Mark van Bakel - Islands Trust